

On the use of a non-extensive distribution in the solar plasma

A. Lavagno and P. Quarati

*Dipartimento di Fisica, Politecnico di Torino, C.so Duca degli Abruzzi 24, I-10129 Torino, Italy
Istituto Nazionale di Fisica Nucleare, Sezioni di Torino e di Cagliari*

Abstract

We describe the physical motivations why we are urged to look for an ionic equilibrium distribution function slightly different from the Maxwellian to calculate nuclear reaction rates in the solar core plasma answering, at the same time, to a recent paper by Bahcall et al. (astro-ph/0010055).

In a recent paper, Bahcall et al. [1] discuss motivations why Salpeter formula for screening of nuclear reactions in the Sun is correct (in contrast with conclusions contained in several other papers) and show five different derivations of the Salpeter screening factor f_0 .

A paper of us [2] is also included in this critical discussion (although our work is not concerned directly on the factorizable screening factor f_0 but, rather, on the evaluation of the reaction rates) because of the use we make of an ionic equilibrium distribution which differs very slightly from the Maxwellian distribution. The reason of using this equilibrium distribution is related to the need of including many-body effects. It was already proposed *ad hoc* by Clayton [3] and we can see, a posteriori, that coincides to the equilibrium distribution of the non-extensive thermostatistics developed by Tsallis [4] and used recently in many different physical applications [5], as for instance, gravitational [6] and high energy problems [7].

In our calculations the screening factor f_0 is taken to be unity because its effect on the reaction rates is negligible if compared to the important depletions (or enhancements) of the standard rates due to other effects (electromagnetic fluctuations in a plasma) responsible of the non-extensive equilibrium distribution function. Also, by following the derivation of f_0 in Ref. [8] and using non-extensive distribution, we can calculate for the solar core about the same value of the screening factor. For comments based on Molecular Dynamics Model (MDM) on the validity of Salpeter screening formula see Ref. [9]. Several authors are actually working on MDM within non-extensive Tsallis statistics [10]. It seems to us obvious that, if all the derivations of the Salpeter formula are based on the validity of the assumptions needed for the Boltzmann-Gibbs statistics, the equilibrium distribution must be, of course, the Maxwellian distribution. However the motivations that deviations from these assumptions are not correct should be explicitly given.

Results within non-extensive statistics go toward a better agreement with the experimental results of the calculated neutrino fluxes. In spite of this achievement, we never excluded the possibility of neutrino oscillation or new physics because only experiments will give the correct answer on this problem. At the same time, we have also shown that modifications

of the reaction rates do not affect bulk properties of the gravitationally stabilized solar core such as sound speed or hydrostatic equilibrium, depending on mean values obtained averaging over Maxwellian distribution function [11].

We confirm that Tsallis distribution is an equilibrium distribution and that the argument that there is enough time to arrive at equilibrium after a reaction is fully accomplished by the Tsallis equilibrium distribution that can be derived in, at least, three different ways containing not alternative reasons.

In one approach we exploit the knowledge that the distribution in the solar interior cannot be too much different from the Maxwellian one and add small corrections (higher-order terms in a derivative expansion) to the coefficients of the standard Fokker-Plank equation. Tsallis' distribution is immediately generated [12].

A second approach focuses on the electric microfields that have been shown to exist in the solar plasma: the time-spatial fluctuations in the particles positions produce specific fluctuations of microscopic electric field in a given point of the plasma. We have shown that such electric microfields in the solar plasma (with energy density of the order of 10^{-16} MeV/fm³) implies a deviation from the Maxwellian distribution; the entity depending on the value of the plasma parameter and on the correlation among ions [2,13].

The third approach has been just started and aims to connect the distribution of collective variables to memory effects and long-time correlations between velocities [14,15]. There should exist solutions compatible with the Tsallis' distribution and/or other non-Maxwellian distributions.

These three approaches are not exhaustive and not necessary alternative. Nevertheless, it is suggestive that all of them point in the same direction: *the Maxwell-Boltzmann distribution of velocity should have small but nonnegligible corrections in the solar plasma and the Tsallis' distribution could provide a better description.*

Normal stellar matter, such as the one in the Sun, is non-degenerate, *i.e.*, quantum effects are small (in fact, they are small for electrons and completely negligible for ions), it is non-relativistic, and it is in good thermodynamical equilibrium. On this ground, the particle velocity distribution is almost universally taken to be a Maxwell-Boltzmann distribution, without much questioning.

However, derivations of the ubiquitous Maxwell-Boltzmann distribution are based on several assumptions [12]. In a kinetical approach, one assumes (1) that the collision time be much smaller than the mean time between collisions, (2) that the interaction be sufficiently local, (3) that the velocities of two particles at the same point are not correlated (Boltzmann's Stosszahlansatz), and (4) that energy is locally conserved when using only the degrees of freedom of the colliding particles (no significant amount of energy is transferred to collective variables and fields). In the equilibrium statistical mechanics approach, one uses the assumption that the velocity probabilities of different particles are independent, corresponding to (3), and that the total energy of the system could be expressed as the sum of a term quadratic in the momentum of the particle and independent of the other variables, and a term independent of momentum, but if (1) and (2) are not valid the resulting effective two-body interaction is not local and depends on the momentum and energy of the particles. Finally, even when the one-particle distribution is Maxwellian, additional assumptions about correlations between particles are necessary to deduce that the relative-velocity distribution, which is the relevant quantity for rate calculations, is also Maxwellian.

At least in one limit the Maxwell-Boltzmann distribution can be rigorously derived: systems that are dilute in the appropriate variables, whose residual interaction is small compared to the one-body energies. In spite of the fact that the effects of the residual interaction cannot be neglected, as a good first approximation the solar interior can be studied in this dilute limit; therefore, it is reasonable to suppose that the velocity distribution in the Sun is not too far from the Maxwellian one.

Likewise, one often assumes that the solar core could be treated as an ideal (Debye) plasma. However, there are physical conditions and/or specific applications that needs higher accuracy for which becomes necessary to take into account modifications of the standard plasma theory.

When $\Gamma \approx 0.05 \div 0.1$, the mean Coulomb energy potential is not much smaller of the thermal kinetic energy and the screening length $R_D \approx a$. It is not possible to clearly separate individual and collective degrees of freedom. The presence of at least two different scales of energies of the same rough size produces deviations from the standard statistics which describe the system in terms of a single scale, kT .

The reaction time necessary to build up screening after a hard collision can be estimated from the inverse solar plasma frequency $t_{pl} = \omega_{pl}^{-1} = \sqrt{m/(4\pi ne^2)} \approx 10^{-17}$ sec, and it is comparable to the collision time $t_{coll} = \langle \sigma v n \rangle^{-1} \approx 10^{-17}$ sec. Therefore, several collisions are likely necessary before the particle loses memory of the initial state and the scattering process can not be considered Markovian. In addition, screening starts to become dynamical: the time necessary to build up again the screening after hard collisions is not negligible any more.

Many remarks we have just made are reported in well known books on statistical mechanics, kinetic and plasma theory and in works by astrophysicist as, for instance, Paquette et al. [16] and Cox et al. [17]. Let us report, specifically, from Cox et al.: *“one usually assumes (1) negligible radiative forces, (2) complete ionization, (3) Maxwellian velocity distributions and the same kinetic temperature for all ions and electrons, (4) diffusion velocities which are much smaller than mean thermal velocities, (5) no magnetic fields, (6) collisions dominated by “classical” interactions between two point particles, and (7) plasma which can be considered a dilute gas, i.e., the ideal gas equation of state applies. [...] When the number of particles in a Debye sphere around an ion is not $\gg 1$, assumption (6) and (7) above are inappropriate. Ions are no longer effectively screened from their surrounding, and multiple particle collisions and collective effects become important. The “classical” approach to calculating transport properties of ions using Boltzmann’s equation becomes invalid. In the Sun, at number for particles per Debye sphere is only about 1.4 at the base of the convection zone, and 2.2 at the core, so the “dilute gas” approximation fails.”*

Finally, let us recall that the introduced equilibrium distribution contains the factor $e^{-(1-q)/2(E/kT)^2}$ where q is the Tsallis parameter. In the solar core plasma we have found for it the value $|(1-q)/2| \approx 0.003$ ($q = 1$ corresponds to the Boltzmann-Gibbs statistics and Maxwellian distribution). In spite of the very small deviation of the behavior at high energy of the Tsallis distribution respect to the standard curve the effect on the reaction rates (depletion or enhancement) is quite important [2,12,13].

REFERENCES

- [1] J.N. Bahcall, S. Brown, A. Gruzinov, R.F. Sawyer, astro-ph/0010055.
- [2] A. Lavagno, P. Quarati, Nucl. Phys. B (PS) 87 (2000) 209.
- [3] D.D. Clayton et al., ApJ. 199 (1975) 494.
- [4] C. Tsallis, J. Stat. Phys. 52 (1988) 479.
- [5] See <http://tsallis.cat.cbpf.br/biblio.htm> for a regularly updated bibliography on the subject.
- [6] A. Lavagno, G. Kaniadakis, M. Rego Monteiro, P. Quarati, C. Tsallis, Astrophys. Lett. Comm. 35 (1998) 449.
- [7] W.M. Alberico, A. Lavagno, P. Quarati, Eur. Phys. J. C 12 (2000) 499.
- [8] J.N. Bahcall, X. Chen, M. Kamionkowski, Phys. Rev. C 57 (1998) 2756.
- [9] G. Shaviv, astro-ph/0010152.
- [10] C. Tsallis, cond-mat/0010150.
- [11] S. Degl'Innocenti et al., Phys. Lett B 449 (1998) 291.
- [12] G. Kaniadakis, A. Lavagno, M. Lissia, and P. Quarati, Physica A 261 (1998) 359.
- [13] A. Lavagno, P. Quarati, preprint Oct. 2000, submitted for publication.
- [14] A. Valuev, A. Kaklyugin, G. Norman, Journ. of Exp. Theor. Phys. 86 (1998) 480.
- [15] M. Coraddu et al., Braz. J. Phys. 29 (1999) 153.
- [16] C. Paquette, C. Pelletier, G. Fontaine and G. Michaud, ApJ. Suppl. 61 (1986) 177.
- [17] A.N. Cox, J.A. Guzik, R.B. Kidman, ApJ. 342 (1989) 1187.